
Professional Certificate in Diabetes Health Coaching Services

Nutrition Therapy for Diabetes

A1C (Hemoglobin A1c) – Related terms: Glycemic control, blood glucose, diabetes monitoring. Explanation: A laboratory measurement that reflects average blood glucose over the past 2-3 months. It is expressed as a percentage; higher values indicate poorer control. Example: An A1C of 7% corresponds roughly to an average glucose of 154mg/dL. Practical application: Use A1C results to set individualized nutrition goals, adjust carbohydrate intake, and evaluate the effectiveness of dietary interventions. Challenges: Variability due to hemoglobinopathies, anemia, or recent blood loss can affect accuracy; therefore, combine with self-monitoring data.

ADA (American Diabetes Association) – Related terms: Clinical practice guidelines, nutrition therapy, diabetes standards. Explanation: Leading professional organization that publishes evidence-based recommendations for diabetes care, including nutrition therapy. Example: The ADA Nutrition Therapy guidelines advise using carbohydrate counting to achieve individualized glycemic targets. Practical application: Reference ADA standards when developing meal plans, counseling on portion sizes, and selecting evidence-based resources. Challenges: Keeping up-to-date with annual guideline revisions and adapting recommendations to diverse cultural food patterns.

AGP (Ambulatory Glucose Profile) – Related terms: Continuous glucose monitoring, glucose variability, time in range. Explanation: A visual summary of CGM data that displays glucose trends over days, highlighting patterns of hyper- and hypoglycemia. Example: An AGP may show that glucose spikes consistently occur after breakfast. Practical application: Use AGP insights to modify breakfast carbohydrate quality, timing, or portion size. Challenges: Requires patient familiarity with CGM technology and interpretation of complex data.

Alkaline diet – Related terms: Acid-base balance, plant-based eating, renal health. Explanation: A dietary pattern emphasizing fruits, vegetables, nuts, and legumes, which are considered “alkaline-forming” after digestion. Example: Incorporating spinach and almond milk can increase the diet’s alkaline load. Practical application: May benefit patients with early diabetic kidney disease by reducing dietary acid load. Challenges: Scientific evidence linking alkalinity to glycemic control is limited; must balance with carbohydrate needs.

Artificial sweetener – Related terms: Non-nutritive sweetener, sugar substitutes, glycemic impact. Explanation: Substances that provide sweetness with little or no calories, such as sucralose, aspartame, and stevia. Example: Adding a packet of sucralose to coffee instead of sugar reduces carbohydrate intake. Practical application: Useful for patients seeking to lower added sugars while preserving taste. Challenges: Some individuals experience gastrointestinal discomfort; long-term effects on gut microbiota remain under investigation.

Basal insulin – Related terms: Long-acting insulin, background insulin, fasting glucose. Explanation: Insulin formulation that provides a steady low-level insulin supply to control glucose between meals and overnight. Example: Insulin glargine is a common basal insulin used once daily. Practical application: Coordinate basal insulin dosing with overall carbohydrate intake to avoid fasting hyperglycemia. Challenges: Matching basal dose to variable daily activity levels and dietary patterns can be complex.

BG monitoring (Blood Glucose monitoring) – Related terms: Fingerstick testing, glucometer, self-monitoring. Explanation: The process of measuring glucose concentration in capillary blood using a handheld device. Example: A patient checks BG before meals, two hours after meals, and at bedtime. Practical application: Provides immediate feedback for adjusting meal composition and timing. Challenges: Patient adherence, cost of test strips, and proper technique affect reliability.

Beta-cell function – Related terms: Insulin secretion, pancreatic health, glucose tolerance. Explanation: The ability of pancreatic beta-cells to produce and release insulin in response to rising glucose. Example: Early type 2 diabetes often shows diminished first-phase insulin secretion. Practical application: Nutritional strategies such as moderate carbohydrate distribution can preserve beta-cell function. Challenges: Progressive loss of beta-cell mass limits the effectiveness of diet alone over time.

Binary diet approach – Related terms: Meal timing, intermittent fasting, carbohydrate cycling. Explanation: A flexible strategy that alternates between days of higher carbohydrate intake and days of lower intake to improve insulin sensitivity. Example: “Carb-rich” days on weekends paired with “low-carb” weekdays. Practical application: May help patients break through plateaus in weight loss or glycemic control. Challenges: Requires careful planning to avoid excessive glucose excursions on high-carb days.

Body mass index (BMI) – Related terms: Obesity, weight status, adiposity index. Explanation: A calculation of weight (kg) divided by height squared (m²) used to categorize underweight, normal weight, overweight, and obesity. Example: A BMI of 32 kg/m² classifies as class I obesity. Practical application: BMI guides calorie targets and the urgency of lifestyle modifications. Challenges: Does not differentiate muscle from fat; may misclassify athletes or older adults.

Bolus insulin – Related terms: Rapid-acting insulin, mealtime insulin, postprandial glucose. Explanation: Short-acting insulin administered to cover the rise in glucose after a meal. Example: Insulin lispro is typically taken within 15 minutes before eating. Practical application: Match bolus dose to carbohydrate amount using an insulin-to-carbohydrate ratio. Challenges: Accurate carb counting and timing are essential; errors can cause hypoglycemia.

Branched-chain amino acids (BCAAs) – Related terms: Protein quality, muscle metabolism, insulin resistance. Explanation: Essential amino acids leucine, isoleucine, and valine, abundant in animal proteins and some plant sources. Example: Whey protein supplements are rich in BCAAs. Practical application: Adequate BCAA intake supports lean-mass maintenance during weight loss. Challenges: Excessive BCAA consumption may exacerbate insulin resistance in some individuals.

Calorie density – Related terms: Energy density, satiety, portion control. Explanation: The number of calories per unit weight (kcal/g) of a food. Example: Fresh fruit has low calorie density compared with chocolate bars. Practical application: Encourage low-density foods to increase volume without excess calories. Challenges: Patients may misinterpret “low-calorie” labels and overconsume.

Carbohydrate counting – Related terms: Carb-to-insulin ratio, glycemic load, meal planning. Explanation: A method of tracking the grams of carbohydrate consumed to match insulin dosing and maintain glycemic control. Example: A 45-g carbohydrate meal may require a bolus of 4 units of rapid-acting insulin if the ratio is 1:10. Practical application: Enables flexible eating patterns while preventing postprandial spikes. Challenges: Requires literacy, numeracy, and consistent food labeling; hidden carbs can be problematic.

Carbohydrate quality – Related terms: Fiber content, glycemic index, whole grains. Explanation: The nutritional value of carbohydrate sources, emphasizing complex carbs with high fiber and low GI. Example: Swapping white rice for quinoa improves carbohydrate quality. Practical application: Improves satiety, reduces postprandial glucose peaks, and supports lipid health. Challenges: Cultural preferences and accessibility may limit options.

Carbohydrate load – Related terms: Carbohydrate density, macronutrient distribution, energy intake. Explanation: The total amount of carbohydrate consumed in a meal or over a day, expressed in grams. Example: A typical “high-carb” dinner may contain 80 g of carbs. Practical application: Adjust load based on activity level and insulin regimen. Challenges: Balancing adequate energy for athletes while preventing hyperglycemia.

Carbohydrate-to-protein ratio – Related terms: Macronutrient balance, glycemic response, satiety. Explanation: The proportion of carbohydrate grams to protein grams in a meal, influencing glucose absorption rate. Example: A 3:1 Carb-to-protein ratio (e.G., 30G carbs and 10g protein) can moderate postprandial spikes. Practical application: Use in mixed-meal planning for people on fixed insulin doses. Challenges: Requires precise food weighing and label reading.

Carotenoids – Related terms: Antioxidants, phytonutrients, eye health. Explanation: Pigments found in colorful fruits and vegetables (e.G., Beta-carotene, lycopene) that have antioxidant properties. Example: Sweet potatoes are high in beta-carotene. Practical application: Encourage carotenoid-rich foods to reduce oxidative stress associated with diabetes complications. Challenges: Fat-soluble nature requires dietary fat for absorption; low-fat meals may limit bioavailability.

Casein protein – Related terms: Dairy protein, slow-digesting protein, muscle synthesis. Explanation: The major protein in milk that digests slowly, providing a steady amino acid release. Example: Cottage cheese provides casein and is often recommended as a bedtime snack. Practical application: Supports overnight muscle repair without causing rapid glucose spikes. Challenges: Lactose intolerance may limit use; portion control is essential due to calorie content.

Celery juice trend – Related terms: Detox diets, low-calorie beverages, fad diets. Explanation: A popular but

scientifically unsupported claim that drinking celery juice improves insulin sensitivity. Example: Some social media posts suggest 16 oz of celery juice each morning. Practical application: Educate patients on evidence-based nutrition; advise moderation and focus on whole foods. Challenges: Overreliance on single-ingredient “cures” can distract from comprehensive therapy.

Cholesterol management – Related terms: LDL, HDL, statins, dietary fat. Explanation: Controlling blood lipids through diet and medication to reduce cardiovascular risk in diabetes. Example: Replacing saturated fat with mono-unsaturated fat can lower LDL cholesterol. Practical application: Integrate cholesterol-friendly foods such as oats, nuts, and fatty fish into meal plans. Challenges: Balancing carbohydrate restriction with heart-healthy fat recommendations.

Combined macronutrient meals – Related terms: Mixed meals, glycemic modulation, satiety. Explanation: Meals that include carbohydrate, protein, and fat in balanced proportions to blunt glucose excursions. Example: A plate with quinoa (carb), grilled salmon (protein/fat), and broccoli (fiber). Practical application: Teach patients to construct meals using the “plate method” to improve glycemic control. Challenges: Portion estimation and cultural food norms may complicate implementation.

Complex carbohydrates – Related terms: Starches, fiber, slow-release carbs. Explanation: Carbohydrate chains that require longer digestion, resulting in gradual glucose release. Example: Whole-grain breads, legumes, and brown rice are complex carbs. Practical application: Prioritize complex carbs to sustain energy and limit spikes. Challenges: Some processed “complex” foods contain added sugars that negate benefits.

Continuous glucose monitoring (CGM) – Related terms: Sensor, glucose trend, time-in-range. Explanation: A device that measures interstitial glucose continuously, providing real-time data and alerts. Example: A CGM may alert a user when glucose falls below 70 mg/dL. Practical application: Use CGM data to fine-tune carbohydrate timing and portion sizes. Challenges: Sensor cost, skin irritation, and data overload for some patients.

Convection cooking – Related terms: Roasting, baking, heat transfer. Explanation: Cooking method where hot air circulates around food, often reducing added fats. Example: Roasting vegetables at 425 °F with a drizzle of olive oil. Practical application: Promote convection methods to lower calorie density while preserving nutrients. Challenges: Requires oven access and familiarity with cooking times.

Co-enzyme Q10 (CoQ10) – Related terms: Mitochondrial antioxidant, supplement, cardiovascular health. Explanation: A lipid-soluble compound involved in cellular energy production; sometimes used to support heart health in diabetics. Example: A daily dose of 100 mg CoQ10 is common in supplement form. Practical application: Consider supplementation for patients on statins who experience muscle pain. Challenges: Limited robust evidence for glycemic impact; interactions with anticoagulants must be monitored.

Coronary artery disease (CAD) – Related terms: Atherosclerosis, cardiovascular risk, lipid profile. Explanation: Narrowing of coronary arteries due to plaque buildup, a leading cause of mortality in diabetes. Example: Patients with A1C >8 % have higher CAD risk. Practical application: Emphasize heart-healthy nutrition (e.G.,

Mediterranean diet) to mitigate CAD development. Challenges: Lifestyle changes may be difficult to sustain without multidisciplinary support.

Cross-contamination (food safety) – Related terms: Gluten-free diet, allergen control, kitchen hygiene. Explanation: Unintended transfer of food particles from one dish to another, potentially altering carbohydrate content. Example: A slice of bread left on a cutting board used for vegetables can add hidden carbs. Practical application: Teach patients to use separate utensils and surfaces when preparing low-carb meals. Challenges: Requires diligent kitchen practices, especially in shared living spaces.

Crude fiber – Related terms: Dietary fiber, insoluble fiber, plant cell walls. Explanation: The portion of plant material that is not digested by human enzymes, contributing to stool bulk. Example: Wheat bran contains high crude fiber. Practical application: Increase crude fiber to improve bowel regularity and modestly attenuate glucose absorption. Challenges: Excessive intake may cause bloating; balance with soluble fiber for optimal effect.

Daily value (DV) – Related terms: Nutrition label, %DV, recommended intake. Explanation: Percentage that indicates how much a nutrient in a serving contributes to the daily recommended amount. Example: A label showing 25 % DV of dietary fiber per serving. Practical application: Use DV to help patients compare foods quickly, especially when scanning packaged items. Challenges: DV is based on a 2,000-calorie diet; individual needs may differ.

Diabetes plate method – Related terms: Visual portion guide, meal composition, carbohydrate allocation. Explanation: A visual tool dividing the plate into halves non-starchy vegetables, a quarter protein, and a quarter carbohydrate. Example: A plate with leafy greens, grilled chicken, and a small serving of quinoa. Practical application: Simplifies portion control for patients without precise measuring tools. Challenges: May not capture specific carbohydrate gram targets needed for insulin dosing.

Diabetes self-management education (DSME) – Related terms: Patient empowerment, behavior change, health coaching. Explanation: Structured educational program that equips individuals with knowledge and skills to manage diabetes. Example: A 6-week group class covering nutrition, physical activity, and medication adherence. Practical application: Integrate nutrition therapy concepts into DSME curricula. Challenges: Attendance barriers, health literacy variability, and resource constraints.

Diabetic ketoacidosis (DKA) – Related terms: Hyperglycemia, ketone bodies, emergency. Explanation: Acute, life-threatening complication characterized by high blood glucose, ketonemia, and metabolic acidosis. Example: A type 1 diabetic who omits insulin may develop DKA. Practical application: Educate patients on carbohydrate-insulin matching to prevent DKA. Challenges: Recognizing early signs and ensuring prompt medical care.

Digestible carbohydrates – Related terms: Simple sugars, maltodextrin, glucose availability. Explanation: Carbohydrates that are readily broken down into glucose and absorbed quickly. Example: Table sugar (sucrose) and honey are highly digestible carbs. Practical application: Use digestible carbs for rapid

correction of hypoglycemia. Challenges: Overconsumption leads to spikes; must differentiate from fiber.

Dietary Approaches to Stop Hypertension (DASH) diet – Related terms: Sodium reduction, potassium, blood pressure. Explanation: Eating pattern rich in fruits, vegetables, whole grains, and low-fat dairy, designed to lower blood pressure. Example: Incorporating berries, leafy greens, and low-fat yogurt daily. Practical application: Beneficial for diabetics with hypertension; aligns with heart-healthy goals. Challenges: Sodium content in processed foods can undermine DASH principles.

Dietary fiber – Related terms: Soluble fiber, insoluble fiber, prebiotic. Explanation: Plant-based carbohydrates resistant to digestion, offering health benefits such as improved glycemic control. Example: Oats contain soluble β -glucan fiber that slows glucose absorption. Practical application: Aim for 25-30 g fiber per day, emphasizing a mix of soluble and insoluble sources. Challenges: Gradual increase needed to avoid gastrointestinal upset; patient misconceptions about “fiber supplements” versus whole foods.

Dietary glycemic index (GI) – Related terms: Carbohydrate quality, glucose response, low-GI foods. Explanation: Ranking of carbohydrate foods based on their impact on postprandial blood glucose compared with a reference (glucose or white bread). Example: A GI of 55 for lentils versus 85 for white rice. Practical application: Pair low-GI carbs with protein/fat to further blunt glucose peaks. Challenges: GI values can vary with ripeness, cooking method, and portion size; not all low-GI foods are low-calorie.

Dietary glycemic load (GL) – Related terms: GI, carbohydrate amount, total impact. Explanation: Product of GI and the amount of carbohydrate in a serving, reflecting overall glycemic effect. Example: A food with GI = 50 and 30 g carbs has GL = 15. Practical application: Use GL to prioritize foods that provide less overall glucose load. Challenges: Requires accurate carb quantification; many food databases lack GL values.

Diuretic effect of high-protein diets – Related terms: Renal load, nitrogen waste, fluid balance. Explanation: High protein intake increases urea production, potentially leading to increased urine output. Example: A 1.5 G/kg protein diet may cause mild diuresis. Practical application: Monitor hydration status in patients on high-protein regimens, especially those on diuretics. Challenges: Over-restriction of fluids can exacerbate kidney strain.

Donor-derived nutrition information – Related terms: Patient-generated data, food diaries, remote coaching. Explanation: Nutrition data collected by patients (e.g., Via apps) shared with health professionals for analysis. Example: A client uploads a week-long food log from a smartphone app. Practical application: Enables personalized feedback without in-person visits. Challenges: Data accuracy depends on patient diligence; privacy concerns must be addressed.

Egg substitution – Related terms: Cholesterol, protein source, plant-based alternatives. Explanation: Replacing whole eggs with egg whites or plant-based equivalents to reduce dietary cholesterol while maintaining protein. Example: Using 2 egg whites instead of 1 whole egg in an omelet. Practical application: Useful for patients with hyperlipidemia who also need protein. Challenges: Loss of yolk nutrients (vitamin D, choline) must be compensated.

Enteral nutrition – Related terms: Tube feeding, formula, malnutrition. Explanation: Delivery of nutrients directly to the gastrointestinal tract via a feeding tube. Example: A diabetic patient with dysphagia receives a diabetes-specific formula. Practical application: Choose formulas with controlled carbohydrate content and added fiber. Challenges: Monitoring glucose levels is critical; formula composition must match insulin regimen.

Essential fatty acids (EFAs) – Related terms: Omega-3, omega-6, inflammation. Explanation: Polyunsaturated fats that the body cannot synthesize and must be obtained from diet. Example: Alpha-linolenic acid from flaxseed is an omega-3 EFA. Practical application: Incorporate EFAs to reduce inflammatory markers associated with insulin resistance. Challenges: Balancing omega-3 to omega-6 ratio; excessive omega-6 may counteract benefits.

Excessive postprandial glucose (PPG) spikes – Related terms: Hyperglycemia, carbohydrate surge, insulin response. Explanation: Sharp rise in blood glucose occurring 1-3 hours after eating, often linked to high-glycemic foods. Example: A 90-minute PPG rise of 70 mg/dL after a sugary snack. Practical application: Advise patients to pair carbs with protein/fat, choose low-GI foods, and monitor portion size. Challenges: Individual variability; stress and activity level also influence PPG.

Fasting plasma glucose (FPG) – Related terms: Diagnostic criteria, basal glucose, oral glucose tolerance test. Explanation: Blood glucose measured after an overnight fast, used to diagnose diabetes and prediabetes. Example: An FPG of 130 mg/dL meets diabetes diagnostic threshold. Practical application: Use FPG results to set baseline calorie and carbohydrate targets. Challenges: Stress, illness, or medication can affect fasting values.

Fiber-rich breakfast cereals – Related terms: Soluble fiber, satiety, glycemic control. Explanation: Breakfast grains high in dietary fiber, such as oatmeal or bran flakes. Example: ½ Cup of rolled oats provides ~4 g soluble fiber. Practical application: Recommend fiber-rich cereals to blunt morning glucose rise and promote fullness. Challenges: Some commercial cereals contain added sugars; label reading essential.

Food frequency questionnaire (FFQ) – Related terms: Dietary assessment, recall bias, nutrition research. Explanation: A survey tool that asks respondents how often they consume specific foods over a defined period. Example: An FFQ may ask how many times per week a person eats fried foods. Practical application: Use FFQ data to identify patterns needing modification (e.g., High sugary beverage intake). Challenges: Relies on memory; may underestimate portion sizes.

Food labeling literacy – Related terms: Nutrition facts panel, ingredient list, %DV. Explanation: Ability to interpret packaged food information to make healthier choices. Example: Recognizing that “contains 12 g sugar” on a label indicates added sugars. Practical application: Teach patients to compare products using serving size, total carbs, and fiber. Challenges: Complex label formats and marketing claims can confuse consumers.

Food swapping technique – Related terms: Portion control, healthier alternatives, caloric reduction.

Explanation: Replacing higher-calorie or higher-glycemic foods with lower-calorie equivalents. Example: Swapping white bread for whole-grain pita. Practical application: Provide a list of common swaps to aid gradual dietary change. Challenges: Taste preferences and cultural habits may limit acceptance.

Free-fat meals – Related terms: Low-fat diet, fat-free dairy, satiety. Explanation: Meals prepared without added fats, often relying on lean proteins and carbohydrates. Example: Grilled chicken breast with steamed vegetables without oil. Practical application: Useful for patients needing strict calorie restriction. Challenges: May reduce flavor and satiety, leading to overeating later.

Fried food avoidance – Related terms: Trans-fat, caloric density, cardiovascular risk. Explanation: Limiting consumption of foods cooked in oil at high temperatures, which increase fat content and oxidative compounds. Example: Opting for baked fish instead of fried fish. Practical application: Recommend baking, steaming, or grilling as healthier cooking methods. Challenges: Accessibility of alternatives and cultural preferences for fried dishes.

Gastric emptying rate – Related terms: Postprandial glucose, fiber, meal composition. Explanation: Speed at which stomach contents move into the small intestine, influencing glucose absorption timing. Example: High-fat meals slow gastric emptying, extending glucose rise. Practical application: Adjust macronutrient balance to modulate emptying for better glucose control. Challenges: Individual variability; gastrointestinal disorders can alter rates unpredictably.

Glycemic control – Related terms: A1C, fasting glucose, target range. Explanation: The overall management of blood glucose levels within desired limits to prevent complications. Example: Achieving an A1C below 7% is a common glycemic control goal. Practical application: Combine carbohydrate counting, portion control, and physical activity to maintain control. Challenges: Lifestyle factors, medication adherence, and stress can disrupt control.

Glycemic index (GI) testing – Related terms: In-vivo studies, reference glucose, food ranking. Explanation: Laboratory procedure measuring blood glucose response to a standardized portion of a test food. Example: Testing boiled potatoes yields a GI of 78. Practical application: Use GI data to guide patient food selections. Challenges: GI values are population averages; individual responses may differ.

Glycemic load (GL) calculation – Related terms: Carbohydrate amount, GI, total impact. Explanation: Formula: $GL = (GI \times \text{carbohydrate grams per serving}) / 100$. Example: A food with 30g carbs and GI = 55 gives GL = 16.5. Practical application: Teach patients to compute GL for mixed meals. Challenges: Requires accurate carb quantification; many databases lack GL values.

Glucose-dependent insulintropic peptide (GIP) – Related terms: Incretin hormone, postprandial insulin, gut-derived signals. Explanation: Hormone released from the small intestine that stimulates insulin secretion in response to nutrient intake. Example: GIP levels rise after a mixed meal containing carbs and fat. Practical application: Recognize that high-fat meals may amplify GIP, affecting insulin needs. Challenges: In type 2 diabetes, GIP response is often blunted.

Glucose monitoring technology – Related terms: Glucometer, CGM, data analytics. Explanation: Devices and software used to track blood glucose levels, ranging from simple fingersticks to advanced sensors. Example: A smartphone app syncs with a CGM to display trends. Practical application: Select appropriate technology based on patient tech comfort and insurance coverage. Challenges: Cost, data security, and user training.

Glucose-to-insulin ratio (GIR) – Related terms: Insulin sensitivity, carbohydrate adjustment, therapeutic index. Explanation: Ratio expressing the amount of glucose that can be metabolized per unit of insulin; often used to assess insulin resistance. Example: A GIR of 1:0.5 Indicates that 1 mg/dL glucose requires 0.5 Units insulin. Practical application: Adjust carbohydrate intake or insulin dose based on observed GIR. Challenges: Requires frequent glucose and insulin data; inter-individual variability is high.

Gluten-free diet (GFD) – Related terms: Celiac disease, wheat alternatives, hidden carbs. Explanation: Elimination of gluten-containing grains (wheat, barley, rye) from the diet. Example: Using rice flour instead of wheat flour in baking. Practical application: For patients with celiac disease and diabetes, ensure gluten-free products are also low-glycemic. Challenges: Many gluten-free processed foods are high in refined starches and sugars.

Glycogen storage disease (GSD) – Related terms: Metabolic disorder, hypoglycemia, carbohydrate therapy. Explanation: Genetic conditions affecting glycogen synthesis or breakdown, leading to abnormal glucose regulation. Example: Type I GSD (von Gierke disease) causes severe fasting hypoglycemia. Practical application: Tailor carbohydrate distribution to prevent hypoglycemia; may require continuous carbohydrate intake. Challenges: Rare condition; specialized nutrition plans needed.

Glycated proteins – Related terms: HbA1c, advanced glycation end-products (AGEs), biomarkers. Explanation: Proteins that have glucose molecules attached, serving as indicators of long-term glucose exposure. Example: Hemoglobin becomes glycated to form HbA1c. Practical application: Use glycated protein levels to monitor chronic glycemic exposure and adjust diet. Challenges: Non-glucose factors (e.g., Anemia) can influence measurements.

Glycemic variability – Related terms: Glucose swings, time-in-range, oxidative stress. Explanation: Fluctuations in blood glucose throughout the day, independent of average levels. Example: Frequent excursions between 60 mg/dL and 180 mg/dL denote high variability. Practical application: Aim to reduce variability through consistent carbohydrate timing and balanced meals. Challenges: Lifestyle stressors and erratic eating patterns increase variability.

Gravimetric portioning – Related terms: Food weighing, kitchen scale, precision. Explanation: Measuring food quantities by weight rather than volume for accuracy. Example: Weighing 120g of cooked quinoa instead of estimating a cup. Practical application: Improves carb counting precision, especially for mixed dishes. Challenges: Requires a scale and willingness to weigh foods at each meal.

Green tea catechins – Related terms: Polyphenols, antioxidant, metabolic rate. Explanation: Bioactive compounds in green tea that may modestly improve insulin sensitivity. Example: EGCG (epigallocatechin

gallate) is the most studied catechin. Practical application: Suggest 2–3 cups of unsweetened green tea as part of a low-calorie beverage plan. Challenges: Caffeine content may affect sleep; excessive intake can cause liver enzyme elevation.

High-glycemic foods – Related terms: Rapid glucose rise, simple sugars, GI >70. Explanation: Foods that cause a swift increase in blood glucose after consumption. Example: White bread, sugary cereals, and ripe bananas. Practical application: Limit high-glycemic foods, especially before physical activity or bedtime. Challenges: Many staple foods fall into this category, requiring careful substitution.

High-protein diet – Related terms: Satiety, lean mass preservation, renal considerations. Explanation: Eating pattern where protein provides 25–30% of total calories, often used for weight loss. Example: A 2,000-kcal diet with 150g protein. Practical application: Supports muscle retention during calorie restriction for overweight diabetics. Challenges: Potential increased renal load; must monitor kidney function.

High-potassium foods – Related terms: Potassium-rich, blood pressure, fruit and vegetable sources. Explanation: Foods containing >400mg potassium per serving, beneficial for blood pressure control. Example: Avocado, spinach, and sweet potatoes. Practical application: Encourage inclusion in meals for patients with hypertension, unless contraindicated by renal disease. Challenges: Patients with advanced kidney disease must limit potassium intake.

Hydration status – Related terms: Fluid balance, urine output, electrolyte balance. Explanation: The degree to which the body maintains adequate water levels, influencing glucose regulation. Example: Dehydration can falsely elevate blood glucose readings. Practical application: Advise regular water intake (≈ 2 L/day) and monitor for signs of dehydration. Challenges: Elderly patients may have blunted thirst response.

Insulin sensitivity – Related terms: Insulin resistance, HOMA-IR, metabolic health. Explanation: The efficiency with which cells respond to insulin to uptake glucose. Example: Sedentary lifestyle reduces insulin sensitivity. Practical application: Promote regular aerobic exercise and balanced macronutrients to improve sensitivity. Challenges: Genetics, obesity, and medications can impair sensitivity despite lifestyle changes.

Insulin-to-carbohydrate ratio (ICR) – Related terms: Bolus calculation, carb counting, dosing. Explanation: The proportion of insulin units needed per gram of carbohydrate (e.g., 1 Unit per 10g carbs). Example: A patient with an ICR of 1:12 Would take 5 units for a 60-g carb meal. Practical application: Teach patients to calculate bolus doses using their individualized ICR. Challenges: Ratios may change with weight loss, activity level, or illness.

Insulin-pump therapy – Related terms: Continuous subcutaneous insulin infusion (CSII), basal rate, bolus wizard. Explanation: Device delivering rapid-acting insulin continuously, mimicking basal secretion, with user-programmed boluses for meals. Example: A pump set to deliver 0.8 Units/hour basal insulin. Practical application: Provides flexibility for variable carbohydrate intake and can improve glycemic variability. Challenges: Requires training, regular site changes, and risk of technical failures.

Intermittent fasting (IF) – Related terms: Time-restricted eating, metabolic switch, fasting periods.

Explanation: Eating pattern alternating periods of fasting (e.G., 16 Hours) with feeding windows. Example: 8-Hour eating window from 12 pm to 8 pm. Practical application: May improve insulin sensitivity and aid weight loss when combined with proper nutrition. Challenges: Risk of hypoglycemia for insulin-treated patients; must adjust medication timing.

International Classification of Diseases (ICD-10) – Related terms: Diagnostic coding, billing, diabetes codes.

Explanation: Standardized coding system for diseases, including E11 for type 2 diabetes. Example: Coding a patient visit as E11.9 (Type 2 diabetes without complications). Practical application: Accurate coding supports reimbursement for nutrition counseling services. Challenges: Requires familiarity with coding updates and payer policies.

Iron-fortified cereals – Related terms: Micronutrient enrichment, anemia, added sugars. Explanation:

Breakfast cereals enriched with iron to prevent deficiency, often containing added sugars. Example: A fortified wheat puff cereal may have 15 g sugar per serving. Practical application: Choose iron-fortified options with low added sugars for diabetic patients at risk of anemia. Challenges: Balancing iron needs with glycemic impact.

Iso-caloric substitution – Related terms: Energy balance, macronutrient swap, weight maintenance.

Explanation: Replacing one food with another that provides the same caloric content but different nutrient profile. Example: Swapping a sugary snack for a piece of fruit with similar calories.